



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Several days later, on the Wilmington and Northern tracks near the same place, I picked up a young Grasshopper Sparrow—also dead but without a mark. All three of these birds were in the juvenile plumage and of about the same age—just beginning to use their wings. Does it show lack of intelligence on the part of the young bird? I believe a great many are killed this way every year and I have never found an adult bird that had been killed by a train.

I believe it is due more to their weak flight than to anything else. They get near the tracks and the suction caused by the passing train draws them beneath it, their wings being too weak to resist the strain.—CHRESWELL J. HUNT, *Philadelphia, Pa.*

Some Massachusetts Records of Interest.—During the past year I have noted several birds of some interest in this region.

Late fall migrants, 1904, include:

Empidonax minimus, Chebec, Concord Turnpike, Lexington, Oct. 2, one.

Dendroica maculosa, Magnolia Warbler, Waltham, Oct. 9, one.

Dendroica striata, Blackpoll Warbler, Cambridge, Nov. 7, one.

An interesting winter record is *Euphagus carolinus*, Rusty Blackbird, Concord Turnpike, Concord, Jan. 22, 1905, one.

Early spring migrants, 1905, include:

Butorides virescens, Green Heron, Charles River, Needham, Apr. 9, one.

Zonotrichia leucophrys, White-crowned Sparrow, Hobb's Brook, Lexington, May 4, one.

Coccyzus erythrophthalmus, Black-billed Cuckoo, Rock Meadow, Belmont, May 4, one.

Other records of interest, 1905:

Oidemia deglandi, White-winged Scoter, one male, Arlington, Spy Pond, May 9.

Bartramia longicauda, Bartramian Sandpiper, Ipswich River, North Reading, May 13, one.

Icterus spurius, Orchard Oriole, Ipswich, May 28, one male.

Vireo noveboracensis, White-eyed Vireo, Ipswich River, South Middleton, June 9, one.—ARTHUR C. COMEY, *Cambridge, Mass.*

Notes from Northwestern Connecticut.—*Otocoris alpestris praticola*.—On May 25, 1905, I secured a pair of these birds in a meadow on the crest of a low ridge about eight miles south of the village of Litchfield. They undoubtedly had a nest there, for they both had been seen in the same meadow the preceding day, and the thin skin of the belly of the female indicated that she probably had a brood. Both were very shy, which I have not found to be the case with these birds later in the season. I believe that this is the first breeding record for the State of Connecticut.

Helminthophila peregrina.—It may be of interest to note that the